Test 5B [2020-01-19] Definite integral



INSTRUCTIONS

READ CAREFULLY

- This questionnaire contains 10 problems, each one of them has 5 different answers; only one of those answers is right, the others are wrong.

 If it were not possible to allow extra time for the development of the assignment, the teacher might have cancelled a few questions, in compliance with the dispensations provided in the customised educational plan.
- Each right answer is worth **5 points**, each wrong answer is worth **0 points** and each problem left unanswered is worth **1 point**. If the answers are delivered after the time limit set by the teacher, a penalty of one point for each minute of lateness will be applied.
- For each problem, **blacken the square** corresponding to the answer believed to be right in the grid printed in the back side of this sheet, with a black or dark blue pen or marker.
 - In case of errors, an answer may be cancelled by blackening the circle below (warning: cancelled answers can not be turned back on); thereafter a new answer may be given.
 - Avoid overstepping the borders of the boxes and draw no other marks in the answers' sheet.
- Correction fluid is forbidden, as well as smartphones and other communication devices. Use questions' sheets for drafts and hand them over at the end of the test, toghether with any additional worksheet. Scientific non-CAS pocket calculators are allowed.

Only the answers accurately copied into the answers' grid will be assessed.

A B C D E 1. _ _ _ _ _ _ _ _ _ _ _ _ _ _ 3. 5. 7. 10.



- 1. If $\int_{-2}^{2} (x^7 + k) dx = 16$, then k =
 - $\boxed{\mathsf{A}}$ -12
 - B 0
 - $\begin{bmatrix} \mathsf{C} \end{bmatrix} 4$
 - D 4
 - E 12
- 2.

$$\int_0^8 \frac{dx}{\sqrt{1+x}} =$$

- A 6
- B 4
- D $\frac{3}{2}$
- E 1
- 3.

$$\int_0^{\frac{\pi}{2}} \frac{dx}{\cos^2 x} =$$

- A $\frac{\pi}{4}$
- B Diverges to infinity.
- C 2
- D 1
- E Does not exist.
- 4.

$$\int_{-1}^{1} \frac{dx}{\sqrt[3]{x^2}} =$$

- A Diverges to infinity.
- B 0
- C 2
- D6
- E Does not exist.

- 5. The mean value of \sqrt{x} over the interval $0 \le x \le 2$ is:
 - $\boxed{\mathsf{A}} \ \tfrac{1}{2}\sqrt{2}$
 - $\boxed{\mathsf{B}} \ \tfrac{4}{3}\sqrt{2}$
 - C $\frac{2}{3}\sqrt{2}$

 - E 1
- 6. What is the mean value of $3t^3-t^2$ over the interval $-1 \leq t \leq 2$?
 - $A \frac{7}{2}$
 - $\boxed{\mathsf{B}} \ \frac{11}{4}$
 - **C** 8
 - $\boxed{\mathsf{D}} \ \frac{33}{4}$
 - E 16
- 7. For which value of p does $\int_0^1 \frac{1}{x^p} dx = 1.25$?
 - |A| p = 2.5
 - $\boxed{\mathsf{B}} p = 0.5$
 - |C| p = 0.2
 - $\boxed{\mathsf{D}} p = 2$
 - E None of the others.
- 8. Evaluate the integral:

$$\int_0^1 \frac{x}{x^2 + 9} dx.$$

- A None of the others.
- $\boxed{\mathsf{B}} \ \tfrac{1}{2} \left(\ln 1 \ln 0 \right)$
- $\boxed{\mathsf{C}} \ \tfrac{1}{2} \left(\ln 10 \ln 9 \right)$
- $\boxed{\mathsf{D}} \ln 1 \ln 0$
- $| E | \ln 10 \ln 9$

9. Evaluate:

$$\int_0^1 x e^x dx.$$

- A 0
- lacksquare B e
- C 1
- $\boxed{\mathsf{D}} \ 2e-1$
- E None of the others.

10. Evaluate the integral:

$$\int_0^4 \frac{x}{x+9} dx.$$

- $\boxed{\mathsf{A}} \ 4 26 \ln 3 + 3 \ln 18$
- $\boxed{\mathsf{B}} \ \frac{4}{13}$
- $\boxed{\mathsf{C}} \ \tfrac{1}{9\ln 13} \ln 3$
- $\boxed{ D } 13 + \ln 3 18 \ln 2$
- $\boxed{\mathsf{E}} \ 4 + 18 \ln 3 9 \ln 13$